

The Role of Telehealth in Reducing the Mental Health Burden from COVID-19

Xiaoyun Zhou, MMH¹ Centaine L. Snoswell, PhD, MPH, BPharm,²
Louise E. Harding, DipMLS,¹ Matthew Bambling, PhD,^{1,3}
Sisira Edirippulige, PhD,² Xuejun Bai, PhD,⁴
and Anthony C. Smith, PhD, MEd, BN, RN^{1,5}

¹Centre for Online Health, Centre for Health Services Research;

²Centre for Health Services Research; and ³Faculty of Medicine,
The University of Queensland, Brisbane, Australia.

⁴Academy of Behavior and Psychology, Tianjin Normal University,
Tianjin, China.

⁵Centre for Innovative Medical Technology, University of Southern
Denmark, Odense, Denmark.

Keywords: Telehealth, Telemedicine, Telepsychiatry, mHealth

The psychological impact of the coronavirus disease 2019 (COVID-19) pandemic must be recognized alongside the physical symptoms for all those affected.^{1,2} Telehealth, or more specifically telemental health services, are practically feasible and appropriate for the support of patients, family members, and health service providers during this pandemic.³ As of March 18, 2020, there were >198,000 COVID-19 infections recorded globally, and 7,900 deaths. Psychological symptoms relating to COVID-19 have already been observed on a population level including anxiety-driven panic buying⁴ and paranoia about attending community events. Students, workers, and tourists who have been prevented from accessing their training institutions, workplaces, homes, respectively, are expected to have developed psychological symptoms due to stress and reduced autonomy and concerns about income, job, security, and so on.⁵ The Chinese, Singaporean, and Australian governments have highlighted the psychological side effects of COVID-19, and have voiced concerns regarding the long-term impacts of isolation and that the fear and panic in the community could cause more harm than COVID-19.⁶⁻⁸

In the absence of a medical cure for COVID-19, the global response is a simple public health strategy of isolation for those infected or at risk, reduced social contact to slow the spread of the virus, and simple hygiene such as hand washing to reduce the risk of infection. While the primary intervention of isolation may well achieve its goals, it leads to reduced access to support from family and friends, and degrades

normal social support systems and causes loneliness, and is a risk for worsening anxiety and depressive symptoms.⁹ If left untreated, these psychological symptoms may have long-term health effects on patients and require treatment adding to the cost burden of managing the illness. Clinical and nonclinical staff are also at risk of psychological distress as they are expected to work longer hours with a high risk of exposure to the virus. This may also lead to stress, anxiety, burnout, depressive symptoms, and the need for sick or stress leave, which would have a negative impact on the capacity of the health system to provide services during the crisis.¹⁰

Treatment protocols for people with COVID-19 should address both the physiological and psychological needs of the patients and health service providers. Providing psychological treatment and support may reduce the burden of comorbid mental health conditions and ensure the wellbeing of those affected. Our challenge is to provide mental health services in the context of patient isolation, which highlights the role of telehealth (through videoconference, e-mail, telephone, or smartphone apps). The provision of mental health support (especially through telehealth) will likely help patients maintain psychological well-being and cope with acute and postacute health requirements more favorably.

Examples of and evidence to support the effectiveness of telemental health are fairly diverse, especially in the context of depression,¹¹ anxiety,¹² and PTSD.¹³ Videoconferencing,¹⁴ online forums,¹⁵ smartphone apps,¹⁶ text-messaging,¹⁵ and e-mails¹⁷ have been shown to be useful communication methods for the delivery of mental health services.

China is actively providing various telemental health services during the outbreak of COVID-19. These services are from government and academic agencies and include counseling, supervision, training, as well as psychoeducation through online platforms (e.g., hotline, WeChat, and Tencent QQ).^{18,19} Telemental health services have been prioritized for people at higher risk of exposure to COVID-19, including clinicians on the frontline, patients diagnosed with COVID-19 and their families, policemen, and security guards. Early reports also showed how people in isolation actively sought online support to address mental health needs, which demonstrated both a population interest and acceptance of this medium.²⁰

Additional telehealth services have been previously funded by the Australian Government (Better Access Initiative

Table 1. Examples of Community Online Mental Health Services Available in Australia

TELEMENTAL HEALTH SERVICES	TARGET MENTAL HEALTH PROBLEMS	TELEHEALTH MODALITIES
Betterhelp https://www.betterhelp.com/	Depression, stress, anxiety, anger, grief, etc.	Text messaging, chat, telephone, videoconferencing, etc.
Black Dog Institute https://www.blackdoginstitute.org.au/	Depression, bipolar disorder, anxiety, suicide and self-harm, post-traumatic stress disorder, etc.	Websites, mobile apps
Moodgym https://moodgym.com.au/	Depression and anxiety	Online self-help platform
E-couch https://ecouch.anu.edu.au/welcome	Emotional problems	Online self-help platform
Beyond Blue https://www.beyondblue.org.au/	Anxiety, depression, and suicide	Online forum
Headspace https://headspace.org.au/	Anxiety, depression, eating disorders, trauma, etc.	Online group chat, e-mail, telephone
RUOK https://www.ruok.org.au/	Suicide	Psychoeducation about how to ask each other about ups and downs

program), to address mental health needs of rural and remote patients during emergency situations,²¹ such as long-term drought and bushfires. In response to COVID-19, the Australian Government has responded with additional funded services through the Medicare Benefits Schedule, enabling a greater range of telehealth services to be delivered, including telehealth consultations with general practitioners and specialists. However, the expanded telehealth program is restricted to special needs groups and the wider population does not have access to the program. A major benefit of expanding telehealth, including mental health, with no restrictions would reduce person-to-person contact between health service providers and COVID-19 and reduce the risk of exposure of noninfected but susceptible patients in waiting room areas. To date, most of the Australian Government's focus has been on managing medical needs of people during the epidemic, rather than providing resources to meet short- and long-term mental health implications. An expansion of access to telemental health support services with a focused public education campaign to promote these services would begin to address this need (Table 1).^{22–24}

Communication of all health needs is important when patients are having to be isolated. We support the use of telehealth as a valuable way of supporting both physical and psychosocial needs of all patients irrespective of geographical location. Simple communication methods such as e-mail and text messaging should be used more extensively to share information about symptoms of burnout, depression, anxiety, and PTSD during COVID-19, to offer cognitive and/or relaxation skills to deal with minor symptoms, and to encourage access to online self-help programs. For people with COVID-

19, telehealth can be used to monitor symptoms and also to provide support when needed.

While there is growing awareness of mortality rates associated with COVID-19, we should also be cognizant of the impact on mental health—both on a short- and a long-term basis. Telemental health services are perfectly suited to this pandemic situation—giving people in remote locations access to important services without increasing risk of infection.

Disclaimer

Health System Sustainability disclaimer: available at (<https://www.healthsystemsustainability.com.au/resources/our-publications/acknowledgements-and-disclaimer/>).

Disclosure Statement

No competing financial interests exist.

Funding Information

We acknowledge the support of the NHMRC Partnership Centre for Health Sustainability (Grant ID No.: 910002 administered by the Australian Institute of Health Innovation, Macquarie University).

REFERENCES

- Shigemura J, Ursano RJ, Morganstein JC, Kurosawa M, Benedek DM. Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: Mental health consequences and target populations. *Psychiatry Clin Neurosci* 2020. [Epub ahead of print]; doi.org/10.1111/pcn.12988.
- Minter A. Quarantine for coronavirus has serious side effects. 2020. Available at <https://www.bloomberg.com/opinion/articles/2020-02-27/coronavirus->

- quarantine-raises-mental-health-concerns-for-china (last accessed March 20, 2020).
3. Smith AC, Thomas E, Snoswell CL, Haydon H, Mehrotra A, Clemensen J and Caffery LJ. Telehealth for global emergencies: Implications for coronavirus disease 2019 (COVID-19). *J Telemed Telecare*. **2020**; in press: doi.org/10.1177/1357633X20916567.
 4. Collett M. Coronavirus update: more COVID-19 cases confirmed across Australia as shoppers stock up on toilet paper, groceries. **2020**. Available at <https://www.abc.net.au/news/2020-03-04/coronavirus-live-updates-toilet-paper-panic-buying-limits/12023040> (last accessed March 20, 2020).
 5. Worthington B. Coronavirus pandemic fears prompt Government to activate emergency response and extend travel ban. **2020**. Available at <https://www.abc.net.au/news/2020-02-27/federal-government-coronavirus-pandemic-emergency-plan/12005734> (last accessed March 20, 2020).
 6. Australian Government Department of Health. Coronavirus (COVID-19). **2020**. Available at <https://www.health.gov.au/health-topics/novel-coronavirus-2019-ncov> (last accessed March 20, 2020).
 7. National Health Commission of the People's Republic of China, Ministry of Civil Affairs of the People's Republic of China. Notice on Strengthening the Psychological Assistance and Social Work Services in the Response to the New Coronary Pneumonia Epidemic Disease. **2020**. Available at www.nhc.gov.cn/jkj/s3578/202003/58a40a1597ff4823b04f84b77931b0d7.shtml (last accessed March 20, 2020).
 8. Lai L. Fear and panic can do more harm than the coronavirus, says PM Lee Hsien Loong. **2020**. Available at <https://www.straitstimes.com/singapore/fear-and-panic-can-do-more-harm-than-the-coronavirus-says-pm-lee-hsien-loong>
 9. Smith EM. Ethnic minorities: Life stress, social support, and mental health issues. *Couns Psychol* **1985**;13:537-579.
 10. Maunder RG. Was SARS a mental health catastrophe? *Gen Hosp Psychiatry* **2009**;31:316-317.
 11. Garcia-Lizana F, Munoz-Mayorga I. Telemedicine for depression: A systematic review. *Perspect Psychiatr Care* **2010**;46:119-126.
 12. Rees CS, Maclaine EJAP. A systematic review of videoconference-delivered psychological treatment for anxiety disorders. *Aust Psychol* **2015**;50:259-264.
 13. Turgoose D, Ashwick R, Murphy D. Systematic review of lessons learned from delivering tele-therapy to veterans with post-traumatic stress disorder. *J Telemed Telecare* **2018**;24:575-585.
 14. Backhaus A, Agha Z, Maglione ML, Repp A, Ross B, Zuest D, et al. Videoconferencing psychotherapy: A systematic review. *Psychol Serv* **2012**;9:111-131.
 15. Kauer SD, Mangan C, Sanci L. Do online mental health services improve help-seeking for young people? A systematic review. *J Med Internet Res* **2014**;16:e66.
 16. Kerst A, Zielasek J, Gaebel W. Smartphone applications for depression: A systematic literature review and a survey of health care professionals' attitudes towards their use in clinical practice. *Eur Arch Psychiatry Clin Neurosci* **2019**;1-14.
 17. Torniainen-Holm M, Pankakoski M, Lehto T, Saarela O, Mustonen P, Joutsenniemi K, Suvisaari J. The effectiveness of email-based exercises in promoting psychological wellbeing and healthy lifestyle: a two-year follow-up study. *BMC psychology* **2016**;4:21.
 18. Ministry of Education The People's Republic of China. Mental health service platform provides around-the-clock psychological support during COVID-19 outbreak. **2020**. Available at http://en.moe.gov.cn/news/press_releases/202003/t20200309_429190.html (last accessed March 20, 2020).
 19. Ministry of Education of the People's Republic of China. In response to the epidemic, the Psychology Department of Beijing Normal University has opened a psychological support hotline and online counselling service (in Chinese) **2020** http://www.moe.gov.cn/jyb_xwfb/moe_1946/fj_2020/202001/t20200128_416724.html (last accessed March 20, 2020).
 20. Liu S, Yang L, Zhang C, Xiang Y-T, Liu Z, Hu S, et al. Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry* **2020**.
 21. Wilson A, Moretto N, Langbecker D, Snoswell CL. Use of reimbursed Psychology Videoconference Services in Australia after their introduction: An investigation using administrative data. *Value in health regional issues* **2020**;21:69-73.
 22. Snoswell C, Mehrotra A, Thomas E, Smith K, Haydon H, Caffery L, Smith AC. Making the most of telehealth in COVID-19 responses, and beyond. **2020**. Available at: <https://croakey.org/making-the-most-of-telehealth-in-covid-19-responses-and-beyond/> (last accessed March 8, 2020).
 23. Ekeland AG, Bowes A, Flottorp S. Effectiveness of telemedicine: A systematic review of reviews. *Int J Med Inform* **2010**;79:736-771.
 24. Black Dog Institute. Self-help tools and apps. **2020**. Available at <https://www.blackdoginstitute.org.au/getting-help/self-help-tools-apps> (last accessed March 20, 2020).

Address correspondence to:

Xiaoyun Zhou, MMH

Centre for Online Health

The University of Queensland

Building 33, Ground Floor, Princess Alexandra Hospital

Queensland 4102

Australia

E-mail: xiaoyun.zhou1@uqconnect.edu.au

Received: March 10, 2020

Revised: March 17, 2020

Accepted: March 17, 2020

Online Publication Date: March 23, 2020